

DEFICIENT PLASMAS

Congenital deficient plasmas (Bottles)

Alpha-2-antiplasmin human deficient plasma (congenital)



Associated products

Human plasma deficient in alpha-2-antiplasmin (acquired)

Informations

α -2-antiplasmin is an inhibitor of serine proteases, mainly plasmin. It plays an important role in the regulation of fibrinolysis.

It has 3 main functions: it inhibits plasmin, interferes with the adsorption of plasminogen to fibrin and binds to the α chain of fibrin.

A decrease in the amount of α -2-antiplasmin can lead to bleeding syndromes.

Reference	Presentation	Format
6-PPDA2APC	Vial	Minimum 50 mL

Plasma from a human donor with congenital α -2-antiplasmin deficiency.

Packaging in bottle. The minimum packaged volume is 50 mL. The price offer is based on the volume requested.

Advantages

Minimize test time.
Ready to use.

Characteristics

Special plasmas are derived from patients with a congenital deficiency, severe or moderate, or presenting a particular profile.
No buffer or preservatives are added. Quickly frozen at -80°C , the plasma maintains perfectly intact the matrix.
All plasmas are stable when stored at -40°C to -80°C .
We carefully pack with dry ice during shipment.
No additive or preservative.
Expiry date > 1 year.
Plastic vials.

