

DEFICIENT PLASMAS

Acquired deficient plasmas (Bottles)

Protein C human deficient plasma (acquired)



Associated products

APC Resistance Kit

C Diluent / S Diluent

CRYOcheck™ Clot C™

Reference

6-PPDPCA

Presentation

Vial

Format

Minimum 50 mL

Packaging in bottle. The minimum packaged volume is 50 mL. The price offer is based on the volume requested.

Informations

Protein C is a 62 kDa glycoprotein, synthesized by the liver in the presence of vitamin K. CP is at the center of a physiological system that inhibits coagulation: the anticoagulant system of protein C. Thrombin associated with thrombomodulin loses its procoagulant properties at the same time as it activates PC to active protein C (PCa).

PCa in the presence of protein S, calcium and phospholipids is capable of cleaving FVa and FVIIIa blocking the amplification loop of thrombin generation.

Advantages

Minimize test time.
Ready to use.

Characteristics

Special plasmas are derived from patients with acquired deficiency, severe or moderate, or presenting a particular profile.
No buffer or preservatives are added.
Quickly frozen at -80° C, the plasma maintains perfectly intact the matrix.
All plasmas are stable when stored at -40° C to -80° C.
We carefully pack with dry ice during shipment.
No additive or preservative.
Expiry date > 1 year.
Plastic vials.

