

Human Gla-domainless Factor X



Associated products

Bovine Factor X
Human Factor X
Mouse Factor X

Informations

A proenzyme or zymogen is a protein precursor of an enzyme which can give, after activation, an active enzyme. Factor X (FX) is a glycoprotein synthesized by the liver, dependent on vitamin K. FX is involved in the common pathway of coagulation. It is activated in FXa by the FT-FVIIa complex or by the FVIIIa-FIXa complex in the presence of phospholipids. FXa is neutralized by TFPI and antithrombin. Gla domains serve to bind calcium ions by chelating them between 2 carboxylic acid residues.

Reference	Presentation	Format
9-HCX-GD	Vial	100 µg
9-HCX-GD-1	Vial	1 mg



Advantages

The vast majority of zymogens is pure (without additives) with > 95 % purity SDS-PAGE. No additive or preservative.

Characteristics

All proteins are accompanied by product information sheets which describe proper storage conditions. In order that we may warrant product stability, it is imperative that these storage conditions be maintained at all times. Many of our protein preparations are formulated in 50 % (vol / vol) glycerol/H₂O which will remain in fluid phase during storage at -20° C. This preferred method of storage yields the greatest protein stability while still allowing access to the stock protein sample without repeated thawing and freezing steps. All products which are formulated with either glycerol/H₂O or aqueous buffer are delivered in microcentrifuge tubes. By briefly centrifuging the samples in their original containers, complete recovery of the sample at the bottom of the tube will be accomplished. Temperatures lower than -30° C should be avoided in order to prevent a phase transition. When preparing to make a dilution of the stock sample, remove the sample from storage at -20° C and place on ice for a brief period of time (5-10 min). The sample will become less viscous and thus easier to pipette. Never allow protein solutions to remain at room temperature for excessive periods of time. Elevated temperatures may enhance the rate of protein degradation. Avoid storing or maintaining dilute protein samples for a long period of time. In general, purified proteins are inherently more stable in concentrated form. Many proteins are «sticky» by nature. To avoid losing protein due to adsorption, extremely dilute protein samples should be prepared in buffers containing excipients such as bovine serum albumin, polyethylene glycol, or gelatin.