LYOPHILIZED CONGENITAL **DEFICIENT PLASMAS**

INTRINSIC PATHWAY FACTOR IX

NATIVE DEFICIENT PLASMAS

Lyophilized plasmas

Factor IX Deficient Plasma Native



















Associated products

Coagulation Control A	
Coagulation Control N	
Coagulation Reference	

Informations

FIX is a vitamin K dependent glycoprotein synthesized by the liver. FIX can be activated to FIXa by FXIa or by FVIIa in the presence of phospholipids and calcium.

A person who is deficient in FIX has hemophilia B.

Reference Presentation Format Number of tests Vial 4-5164008 5 x 1.0 mL 100 Vial 4-5164016 50 x 1.0 mL 1000

Plasma deficient for Factor IX assay.

Native Factor IX Deficient Plasma is a freeze-dried, native (congenital deficiency) hemophilia B donor plasma pool with a coagulant activity <1% for FIX. All other coagulation factors have normal values.

Components

- 5 or 50 vials x 1 mL lyophilized plasma

Characteristics

- Deficient human plasma, stabilized and lyophilized with an activity <1% of the corresponding coagulation factor.
- All other coagulation factors have normal values.
- Native FIX deficient plasma is used for the determination of FIX by a one-step method based on Cephalin-Kaolin Time (TCK).
- Does not contain inhibitors



