DEFICIENT PLASMAS

Immunodepleted deficient plasmas

Fresh frozen plasmas

Plasminogen Immunodepleted Deficient Human Plasma











Associated products

a2-Antiplasmin Immunodepleted Deficient Human

Antithrombin deficient plasma immuno depleted Antithrombin Immunodepleted Deficient Human Plasma

Reference	Presentation	Format
6-FDPPLG	Bottle	1 x 100 mL
6-FDPPLG-10	Kit	10 x 1.0 mL

Immunodepleted deficient plasma deficient for plasminogen assay

Pooled normal citrated human plasma depleted of plasminogen using antibodies directed to plaminogen immobilized on agarose beads. Plasma contains 20 mM Hepes.

Informations

Plasminogen is the zymogen of plasmin, a key enzyme in the fibrinolysis system. Plasminogen is synthesized mainly by the liver but also the eosinophils, the kidney and the cornea.

It exists in 2 molecular forms: glu-plasminogen (native form) and lys-plasminogen (more active form). The main pathways for activating plasminogen to plasmin involve t-PA and u-PA.

Components

- 10 cryotubes x 1 mL or 100 mL vial of frozen plasma.

Advantages

- No bovine additives
- No reconstitution error
- No deterioration of plasmas linked to freeze-drying
- Cryotubes ready to use after thawing (4 min at 37°C).

Characteristics

Packaging in plastic cryotubes or in bottles of at least 100 mL.

Frozen, immuno-depleted plasmas are certified to have less than 1% for the deficient factor considered, both for the antigenic and functional assay in haemostasis.

This box is intended for research use

